

Measuring Neutron Star Mass and Radius with Three Mass-Radius Relations

C. M. Zhang¹, H.X. Yin¹, Y. Kojima², H. K. Chang³, R. X. Xu⁴, X. D. Li⁵, B. Zhang⁶, B. Kizilta

1. National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

2. Department of Physics, Hiroshima University, Higashi-Hiroshima 739-8526, Japan

3. Department of Physics and Institute of Astronomy, National Tsing Hua University, Hsinchu 30013, Taiwan, China

4. Department of Astronomy, Peking University, Beijing 100871, China

5. Department of Astronomy, Nanjing University, Nanjing 210093, China

6. Department of Physics, University of Nevada, NV 89154-4002, USA

7. Department of Astronomy & Astrophysics, University of California, Santa Cruz, USA

3 February 2008

ABSTRACT

We propose to determine the mass and the radius of a neutron star (NS) using three measurable mass-radius relationships, namely the “apparent” radius inferred from neutron star thermal emission, the gravitational redshift inferred from the absorption lines, as well as the averaged stellar mass density inferred from the orbital Keplerian frequency derived from the kilohertz quasi periodic oscillation (kHz QPO) data. We apply the method to constrain the NS mass and the radius of the X-ray sources, 1E 1207.4-5209, Aql X-1 and EXO 0748-676.

Key words: star: neutron — X-ray: stars — equation of state — pulsar: general

1 INTRODUCTION

Measuring the mass (M) and the radius R of a neutron star (NS) is very important in both nuclear physics and gravitational physics since it allows us to constrain the NS matter compositions (e.g. neutrons or quarks) at super-density (see, e.g. Lattimer & Prakash 2004; Miller 2002; Li et al. 1999; Cheng et al. 1998; Haensel et al. 1986; Alcock et al. 1986), and to trace the particle motion behavior in super-strong gravitational field (see, e.g. van der Klis 2000, 2006).

NS masses have been measured in the radio pulsar binary systems with high precision (see, e.g. Kaspi et al. 1994; Bailes et al. 2003). For example, the masses of the two NSs of the Hulse-Taylor pulsar system, PSR 1913+16, have been measured as $M=1.41M_{\odot}$ and $M=1.38M_{\odot}$, respectively (see, e.g., Manchester & Taylor 1977); and those of the recently discovered double pulsar system, PSR J0737-3039, are $M=1.34M_{\odot}$ and $M=1.25M_{\odot}$, respectively (see, e.g., Lyne et al. 2004). The mass constraints of 26 neutron stars in the binary radio pulsar systems have been derived by Thorsett & Chakrabarty (1999), who present a remarkably narrow underlying gaussian mass distribution at $M=1.35\pm0.04M_{\odot}$. In the low mass X-ray binaries (LMXBs) a lower mass limit $M > 0.97 \pm 0.24M_{\odot}$ was derived for the NS in 2A 1822-371 (Jonker et al 2003), and the NS mass in Cyg X-2 was derived as $M = 1.78 \pm 0.23M_{\odot}$ (Orosz & Kuulkers 1999).

In contrast, there is no method so far to directly measure the radius of a NS. A conventional value of NS radius is about 15 km, which is derived by requiring that the averaged dipolar magnetic field strength (i.e. $\sim 10^{12}$ G) inferred from radio pulsar spindown

(see, e.g. Manchester & Taylor 1977) is consistent with the one implied by the cyclotron line emission in some X-ray binaries (see, e.g. Makishima et al. 1999; Truemper et al. 1978). The uncertainties in determining NS radius make it difficult to accurately measure the NS equation-of-state (EOS). The detail of matter composition inside a NS has been an open issue.

With the launches of X-ray space observatories, especially the recent XMM-Newton, Chandra, and RXTE (see, e.g. Jansen et al. 2001; van der Klis 2000, 2006), some $M - R$ relationships have been appropriately measured or estimated. These include (1) the “apparent radius” R_{∞} estimated from the thermal emission of isolated NSs (e.g. Geminga, Caraveo et al. 2004; 1E 1207.4-5209, Bignami et al. 2004; RX J1856.5-3754, Truemper et al. 2004; Burwitz et al. 2003; Burwitz et al. 2001) or NSs in binary systems (e.g. Aql X-1, Rudledge et al. 2001); (2) the gravitational redshift implied from the absorption lines in EXO 0748-676 and in 4U 1700+24 (e.g., Cottam et al. 2002; Tiengo et al 2005) and (3) the averaged NS mass density inferred from the orbital Keplerian frequency used to interpret the kHz QPOs discovered by RXTE (see, e.g., Miller et al. 1998; van der Klis 2000, 2006; Zhang 2004).

Generally speaking, if two of the three above $M - R$ relations are well “measured”, one can solve both M and R independently. This is the main topic of this paper. We will discuss the methods in detail and apply them to several sources. **Once NS M and R known, we can evaluate the matter compositions of star by comparing with the representative EOSs, as shown in the three figures we chosen the EOSs of strange matter (CS1 and CS2), normal neutron matter (CN1 and CN2) and pion conden-**

sate in the star core (CPC) (see, e.g. Lattimer & Prakash 2001; Cook et al. 1994).

2 MEASURING NS MASS AND RADIUS WITH $M - R$ RELATIONS

(1) The “apparent radius” R_∞ inferred from the thermal emission data is only an upper limit of the NS radius. It is related to the “true” radius R through (see, e.g., Haensel 2001; Lattimer & Prakash 2001; Thorne 1977),

$$R_\infty = R / \sqrt{1 - R_s/R}, \quad (1)$$

where $R_s = 2GM$ is the Schwarzschild radius. From Eq.(1), the maximum value of the mass can be easily obtained as $M = 1.3(R_\infty/10^6 \text{ cm})M_\odot$ when $R = 0.58R_\infty$. **We stress that the apparent radius is not a directly observable quantity although derivable (in principle) from two observable quantities, namely the “apparent blackbody luminosity” (the blackbody luminosity seen by a distant observer) and the “apparent blackbody temperature”. Both quantities are derived by fitting X-ray data under the assumption of perfect blackbody, which is, in general, not true (see, Haensel 2001). In particular the inclusion of an atmosphere above the NS crust can vary the estimate of the NS apparent radius by more than a factor 2 (see Haensel 2001, section 3.1), thus the ‘measured’ apparent radius is still model dependent.**

(2) The gravitational redshift near the neutron star surface provides another M-R relation, giving the ratio between the NS mass and the NS radius (e.g. Miller 2002). The approach becomes relevant with the recent discovery of the absorption features in EXO 0748-676 (Cottam et al. 2002). According to the definition of the gravitation redshift $z = \Delta\lambda/\lambda \simeq GM/R$ (for a weak gravitational field, where λ is the wavelength of the emission), for a non-rotating spherical gravitational source in the Schwarzschild spacetime, we get (see, e.g., Sanwal et al. 2002),

$$1 + z = 1 / \sqrt{1 - R_s/R}, \quad (2)$$

or, in terms of M/R ,

$$m/R_6 = f(z) = \frac{z(1 + z/2)}{0.15(1 + z)^2}, \quad (3)$$

where $R_6 = R/10^6 \text{ cm}$, and $m = M/M_\odot$ is the mass in unit of solar mass. In the case of weak gravitational fields ($z \ll 1$), $f(z) \simeq z/0.15$ is approximately obtained.

(3) A third $M - R$ relation can be obtained by analyzing the kilohertz quasi periodic oscillation (kHz QPO) data. This field has been greatly benefited from the RXTE mission, which discovered the twin (upper and lower) kHz QPOs or millisecond variability in the X-ray data of about 20 NS LMXB systems. The upper kHz QPO frequency is generally interpreted as the Keplerian frequency ν_K of orbital materials at some preferred radius r near the NS surface (van der Klis 2000, 2006; Morsink 2000; Stella & Vietri 1999; Zhang 2004), i.e.,

$$\nu_K = \sqrt{\frac{GM}{4\pi^2 r^3}} = 1850 \text{ (Hz)} A X^{3/2}, \quad X = R/r, \quad (4)$$

with the parameter $A = (m/R_6^3)^{1/2}$ or $R_6 = 1.27 m^{1/3} (A/0.7)^{-2/3}$, where R_6 is the NS radius in unit of 10 km. The physical meaning of the parameter A is such

that the quantity A^2 represents the “measurement” of the averaged NS mass density, namely,

$$M/R^3 \simeq 10^{15} (\text{g/cm}^3) (A/0.7)^2. \quad (5)$$

Meanwhile, by interpreting the twin kHz QPOs as the Keplerian motion and the Alfvén wave oscillation mechanism (see, e.g., Zhang 2004), the averaged mass density parameter A can be inferred from the simultaneously detected twin kHz QPO frequencies. For typical kHz QPO sources, $A \simeq 0.7$ is approximately obtained. If, instead, only one single kHz QPO frequency is detected, one can not derive the exact value of A from the model, but can estimate the lower limit of A and the upper limit of M according to the following arguments, as proposed by Miller et al. (1998). In order to explain the observed X-ray flux saturation at the twin kHz QPO frequencies (e.g. van der Klis 2000, 2006; Zhang et al. 1998), it is usually believed that the maximum Keplerian frequency ν_K occurs either at the star surface ($r = R$: $X = 1$) or at the innermost stable circular orbit (ISCO) with the radius $R_{\text{ISCO}} = 3R_s = 6GM$. The two inequalities are therefore derived from Eq.(4) of Miller et al (1998), i.e.

$$A \geq \nu_K / 1850 (\text{Hz}), \quad (6)$$

and

$$m \leq 2200 (\text{Hz}) / \nu_K. \quad (7)$$

In addition, Burderi & King (1998) have derived a $M - R$ relation for the millisecond accretion-powered X-ray pulsar SAX J 1808.4-3658 with the measured spin frequency 401 Hz through comparing the stellar radius with its co-rotation radius of the magnetosphere-disk. They inferred a value of $A > 0.62$.

To summarize, hitherto there are roughly three kinds of well-known $M - R$ relationships that could be measured or estimated from the data, as described above. Knowing two of the three in a system would in principle lead to the determination of both M and R . In reality, errors are large for these relationships. Knowing all three relations for a source would greatly diminish the errors. As an illustration, below we will use the method to constrain M and R for several sources.

2.1 Constraining M and R with R_∞ and z : the case of 1E 1207.4-5209

If the “apparent radius” R_∞ and the gravitational redshift z are known, both M and R can be calculated by the following two formulae, derived from Eqs.(1) and (2),

$$R_6 = R_\infty / (1 + z), \quad (8)$$

$$m = f(z) R_\infty / (1 + z), \quad (9)$$

with $R_\infty = R_\infty / 10^6 \text{ cm}$.

In the spectral analysis of 1E 1207.4-5209, Bignami et al. (2004) found that the best fitting continuum model includes two blackbody components, i.e. a cooler component with a temperature $KT = 0.163 \pm 0.003 \text{ keV}$ and an emitting radius of $R_e = 4.6 \pm 0.1 \text{ km}$, and a hotter component with $KT = 0.319 \pm 0.002 \text{ keV}$ and $R_e = 0.83 \pm 0.03 \text{ km}$. If we take the cooler component as the emission from the whole star, then the cooler emitting radius $4.6 \pm 0.1 \text{ km}$ can be regarded as a measured lower limit of the “apparent radius” R_∞ . **In addition, an emitting radius $R = 4.6 \pm 0.1 (d/2 \text{ kpc}) \text{ km}$ in 1E 1207.4-5209 is obtained by the best fitting continuum model of the blackbody function and by the distance of 2 kpc**

(Bignami et al. 2004), however as claimed by Haensel (2001) that an emitting radius may be doubled if the atmosphere model of the photon spectrum is considered, i.e. the emitting radius may be as high as ~ 9.4 km.

On the other hand, two absorption features in the spectrum of 1E 1207.4-5209 were detected with the *Chandra* ACIS (Sanwal et al. 2002). Interpreting these lines as Helium atomic lines, a gravitational redshift is inferred as $z \sim (0.12 - 0.23)$ (Sanwal et al. 2002). This conclusion is, however, inconclusive, since the lines may be due to the cyclotron origin (e.g. De Luca et al. 2004). Here we take this gravitational redshift measurement as an unconfirmed, tentative case.

In Figure 1, we present the $M - R$ diagram of NSs with various EOSs and present the above two $M - R$ constraints in the same plot. For the possible parameter ranges of R_∞ and $z = 0.12 - 0.23$, i.e. $R_\infty = 4.5 - 4.7$ (km) as estimated from the blackbody assumption (Bignami et al 2004), $R_\infty = 9.4$ (km) while the atmosphere model of photon spectrum is taken into account (Haensel 2001), and $R_\infty = 20$ (km) assumed under the condition that the uncertainty in the distance of the source 1E1207.4-5209 to the earth is considered (Truemper 2005), we constrain the parameter space within the area enclosed by the lines labeled R1, R2, R3, and those labeled $z=0.12, 0.23$, where the implied $M - R$ ranges are given. In Figure 1, we also plot the lines of $R = R_s = 2GM$ (the solid straight line labeled as R_s , below which the $M - R$ values will be permitted for NSs) and $R = R_{\text{ISCO}} = 6GM$ (the middle straight dotted line labeled as ISCO). The latter corresponds $z = 0.23$ (this value is fortuitous coincidence with the observationally inferred redshift upper limit of 1E1207.4-5209, see Sanwal et al. 2002) and divides the $M - R$ diagram into the $R < R_{\text{ISCO}}$ region (above the line) and the $R > R_{\text{ISCO}}$ region (below the line). From Figure 1 the inferred $M - R$ value regime for 1E1207.4-5209 favors the existence of the exotic matter inside NS. However, it is pointed out by Truemper (2005) that the distance d from 1E 1207.4-5209 to the earth is not so certain and $d = 1.3 \sim 3.9 \text{ kpc}$ has been given (see, e.g. Pavlov et al. 2004). If we consider the atmosphere model of photon spectrum and the distance $d = 3.9 \text{ kpc}$, then the derived “apparent radius” can be as large as 18 km ($4.6 \times 2 \times 3.9/2 = 18$), which will weaken the conclusion of 1E 1207.4-5209 including a strange star because the inferred $M - R$ range in Figure 1 is enlarged to cover the EOSs of normal neutrons.

If one assumes the validity of the above measured R_∞ (4.5 - 4.7 km) and z (0.12 - 0.23), then one gets $M = 0.34 \pm 0.09 M_\odot$ and $R = 4.2 \pm 0.1 \text{ km}$ for 1E 1207.4-5209 according to Eqs.(8) and (9). In this case, one can conclude the NS EOS of 1E 1207.4-5209 infers the composition of strange quark matters (e.g. Xu 2005). Apparently, this conclusion is model dependent and based on the blackbody assumption and the distance of 2 kpc .

2.2 Constraining M and R with R_∞ and A : the case of Aql X-1

If the apparent radius R_∞ and the parameter A are known, both M and R can be calculated by the following two formulae, derived from Eqs.(1) and (5),

$$R_6 = R_{\infty 6} / \sqrt{1 + 0.15(A/0.7)^2 R_{\infty 6}^2}, \quad (10)$$




Figure 1. The $M - R$ diagram. Five representative EOSs are shown (see also, e.g. Miller 2002): stars containing strange quark matter (CS1 and CS2), stars made of normal neutron matter (CN1 and CN2), and stars with pion condensate cores (CPC) (Lattimer & Prakash 2001; Cook et al. 1994). The straight lines are constant z lines: the solid (R_s) and the dotted lines (ISCO) represent that the stellar radius is one and three times of the Schwarzschild radius, i.e. $R = 2GM$ (below the line R_s the NS $M - R$ regimes are possible, see e.g. Haensel et al. 1986) and $R = 6GM$, respectively. The constant redshift lines $z = 0.12$ and $z = 0.23$ are also plotted. The concave-down parabolas are constant R_∞ lines: $R_\infty = 4.5, 9.4, 18 \text{ km}$ labeled by R1, R2 and R3 correspond the lower limit of the apparent radius of 1E1207.4-5290 ($4.5=4.6-0.1$) inferred by the blackbody assumption (Bignami et al. 2004), ($9.4=2 \times (4.6+0.1)$) estimated by taking into account the factor 2 uncertainty because of the inclusion of the NS atmosphere (Haensel 2001)), and by considering the possible longer distance from the source to the earth $d \sim 3.9 \text{ kpc}$ (see, e.g. Pavlov et al 2004), respectively. The shadowed area stands for the possible $M - R$ range of 1E1207.4-5209, which covers the strange matter and the normal neutron EOSs.

$$m = 0.49(A/0.7)^2 \left(R_{\infty 6} / \sqrt{1 + 0.15(A/0.7)^2 R_{\infty 6}^2} \right)^3. \quad (11)$$

The quiescent spectrum of the transient type-I X-ray bursting NS Aql X-1 is measured with *Chandra*/ACIS-S, and the best-fit value of the apparent radius $R_\infty = 13.4^{+5}_{-4} (d/5 \text{ kpc}) \text{ km}$ is inferred by assuming a pure hydrogen atmosphere with the temperatures ranging from 145–168 eV, plus a power-law component. The distance from the observer to the source is 5 kpc as a fiducial value because the current uncertainties allow for a distance from 4 kpc to 6.5 kpc (Rutledge et al. 2001). Therefore, including the uncertainty on source distance the lower and upper limits on R_∞ are 7.5 km ($(13.4 - 4) \times 4/5$) and 23.9 km ($(13.4+5) \times 6.5/5$), respectively. Moreover, the single kHz QPO frequency 1040 Hz has been detected in Aql X-1 by RXTE (e.g. van der Klis 2000, 2006), so that according to Eqs. (6) and (7), one can get the constraints $A \geq 0.56$ and $m \leq 2.1$. In Figure 2, the ranges of M and R of NS in Aql X-1 are confined in the shadowed area where the meanings of the boundaries are indicated in the figure. We find that the shadowed area has a loose constraint on the NS EOSs, and it covers the representative chosen EOSs of the strange matter, the normal neutrons and the pion condensate in the star core. Moreover, it is worth noticing that the $A = 1.0$ parabola divides the $M - R$ diagram




Figure 2. Similar to Figure 1. The mass and radius of NS in Aql X-1 are constrained in the shaded area. R1 (R2) represents $R_\infty = 7.5(23.9)$ (km), where the lower and upper limits of R_∞ on account of the uncertainty by the source distance are 7.5 km $((13.4 - 4) * 4/5)$ and 23.9 km $((13.4+5)*6.5/5)$, respectively. The two concave-up parabolas stand for $A = 0.56$ and $A = 1.0$, respectively (as marked). The horizontal line $m = 2.1$ represents $M=2.1M_\odot$.

into two parts (Figure. 2), and the strange star EOSs are relevant only when $A > 1.0$

2.3 Constraining M and R with A and z : the case of EXO 0748-676

If the gravitational redshift z and the parameter A are known, both M and R can be calculated by the following two formulae, derived from Eqs.(2) and (5),

$$R_6 = 1.43 f^{1/2}(z) (A/0.7)^{-1}, \quad (12)$$

$$m = 1.43 f^{3/2}(z) (A/0.7)^{-1}. \quad (13)$$

In case that only a single kHz QPO frequency is detected, we can place a lower limit on the parameter A from Eq.(6). Following Eqs.(12) and (13), M and R are constrained by the following inequalities

$$R_6 \leq (1850(\text{Hz})/\nu_K) f^{1/2}(z), \quad (14)$$

$$m \leq (1850(\text{Hz})/\nu_K) f^{3/2}(z). \quad (15)$$

A gravitational redshift $z = 0.35$ was detected in EXO 0748-676 by Cottam et al. (2002), who identify the most significant features with the Fe XXVI and XXV n=2-3 and O VIII n=1-2 transitions, all with a redshift of $z = 0.35$, identical within small uncertainties for the respective transitions, however this small error has not been given quantitatively by the authors. In order to present the possible influence of the redshift variation on the $M-R$ region, we plot $z = 0.3$ and $z = 0.4$ lines in Figure 3. A single kHz QPO frequency 695 Hz is detected by RXTE for the same source (e.g., Homan & van der Klis 2000; van der Klis 2000). Thus $A \geq 0.38$ and $m \leq 3.2$ are inferred from Eq.(6) and Eq.(7). As shown in Figure 3, the $A \geq 0.38$ condition gives a very loose constraint, which has little help to constrain the




Figure 3. Similar to Figures 1 and 2. The mass and radius of the NS in EXO 0748-676 are constrained by the line $z = 0.35$ and two horizontal lines representing one (three) solar mass(es). Cottam et al. (2002) declaimed the gravitational redshift $z = 0.35$ with small error, but this small error is not given quantitatively. As a possible comparison, we plot the lines with the redshift $z = 0.3$ and $z=0.4$.

NS mass into a domain of less than $3 M_\odot$ because the detected QPO frequency of this source is abnormally lower than the typical values 1000 Hz (see, van der Klis 2000, 2006). If the future detections of this source present the higher frequency, then the promising constrain condition would be improved. Moreover, since $M=0.97 \pm 0.24M_\odot$ was measured as a lower mass for 2A 1822 - 371 (Jonker et al 2003), we set $1.0 M_\odot$ and $3.0 M_\odot$ as the conservative lower and upper mass limits for NSs, respectively. The $M-R$ constraint for EXO 0748-676 is then limited to the segment of the straight dashed line $z = 0.35$ between the two horizontal lines $m = 1.0$ and $m = 3.0$, where the EOSs for the strange matter, normal neutron and pion condensate in the star core are all permitted.

3 SUMMARY AND DISCUSSION

In this paper, we discuss the possibility of using any two of the three $M-R$ relationships to constrain NS mass and radius. The three $M-R$ relationships include the “apparent radius” (R_∞), the gravitational redshift (z) and the averaged mass density (A^2). We applied the method to three sample sources, 1E1207.4-5209, Aql X-1 and EXO 0748-676. In principle, if all the above three $M-R$ relations are well measured in one source, one can pose tight constraints on the ranges of M and R , so that the potential NS EOSs would be inferred. Unfortunately, this is not achieved so far. Furthermore, some $M-R$ relations are incomplete or have large errors, so that additional astrophysical arguments are needed, and one can derive some inferred lower or upper limits of the $M-R$ relations (e.g. Eqs.[6] and [7]). In binary systems, if the NS mass is known, any one of the three $M-R$ relation measurements would be sufficient to derive its radius.

For the compact star in 1E1207.4-5209, the estimates of its mass and radius come from the possible lower limit of the calculated apparent radius of about 4.6 km (Bignami et al. 2004). An extraordinary low mass is inferred, which is even less than the mea-

sured lower limit of NS mass $0.97 \pm 0.24 M_{\odot}$ in 2A 1822 - 371 (Jonker et al. 2003). In such a case, the existence of exotic matter inside the star is hinted under the assumptions of blackbody spectrum and $2kpc$ distance.

The stellar radius can be also estimated by the motion-induced Doppler-broadening of absorption lines, when the emission region moves towards and away from the observer as the star rotates. According to this, Villarreal and Strohmayer (2004) have studied the source EXO 0748-676 with the measured gravitational redshift $z = 0.35$ (Cottam et al. 2002), and obtained a radius of about 11.5 km and a stellar mass of about $1.75 M_{\odot}$. A remark is that the emission latitude and the orientation of the rotation axis both affect the result, so that the method is model-dependent. In addition, from the recent further study, Chang et al. (2005) show that the intrinsically broad line profile prohibits any meaningful constraint on the NS radius if the 45 Hz burst oscillation seen by Villarreal & Strohmayer (2004) is the spin frequency.

It is worth to mention the measured apparent radius of isolated NS in RX J1856.5-3754 because the distance of this source to earth is well determined to be about 117 pc (see, e.g. Walter & Lattimer 2002), with the perfect black-body spectrum of this source (Drake et al. 2002; Burwitz et al. 2003, 2001) the conservative lower limit of apparent radius has been implied to be 16.5 km ($d/117$ pc) (Truemper et al. 2004). This corresponds to the "true" radius of 14 km for a $1.4 M_{\odot}$ neutron star, indicating a stiff equation of state at high densities, which excludes the quark stars or even the NSs with the quark cores (Truemper et al. 2004). Moreover, without knowledge of the spectrum line in RX J1856.5-3754 until now (see e.g. Drake et al. 2002), we cannot constrain its M and R independently (Truemper et al. 2005). **Once again, it is noted that the above conclusion comes from the black-body spectrum assumption, and the apparent radius may confront a modification by more than a factor of 2 if using the atmosphere model of photon spectrum (Haensel 2001).**

The next generation X-ray observatories with enhanced spectral and timing capabilities would greatly improve the "measurements" of the three $M - R$ relations discussed in this paper. This would strengthen our confidence in evaluating the super-dense nuclear compositions of compact stars. Theoretically, if both M and R of a compact star are accurately measured, its EOS and nuclear matter compositions would be explicitly revealed. This in turn sheds light on the unknown astrophysical processes during the supernova explosion of its progenitor star (e.g. Podsiadlowski et al. 2005). On the other hand, the regime of non-Newtonian strong gravitational fields would be also revealed with the determination of the NS mass and radius, where the direct tests of Einstein's theory of general relativity will be possible in the near future (e.g. van der Klis 2000, 2006).

4 ACKNOWLEDGEMENTS

We are grateful for M.C. Miller for providing the EOS data files. Thanks are also due to the helpful discussions with V. Burwitz, J. Truemper, J.M. Lattimer. We especially thank J. Truemper for many suggestions and advices, and for initialing discussions on $M-R$ relation of RX J1856.5-3754. Thanks are also duo to the helpful discussions with K.S. Cheng, Z.G. Dai, D.M. Wei, T. Lu, Q.H. Peng, Z.R. Wang, J.L. Han, G.J. Qiao, and X.J. Wu. This research has been supported by the innovative project of CAS of China. We are very grateful for the critic comments from the anonymous referee that greatly improve the quality of the paper.

Note added: after acceptance of the paper, we paid attention to the work by F. Ozel on the mass and radius of NS in EXO 0748-676 (Nature 2006, 441, 1115)

REFERENCES

- Alcock, C. et al. 1986, Phys. Rev. Lett., 57, 2088
 Bailes, M., Ord, S.M., Knight, H.S., & Hotan, A.W. 2003, ApJ, 595, L49 (astro-ph/0307468)
 Bignami, G.F., De Luca, A., Caraveo, P.A., Mereghetti, S., Moroni, M., & Mignani, R.P. 2004, Mem. S.A.It, 75, 448 (astro-ph/0402231)
 Burderi, L., & King, A.R. 1998, ApJ, 505, L135
 Burwitz, V., Haberl, F., Neuhaeuser, R., Predehl, P., Truemper, J., & Zavlin, V.E. 2003, A&A, 399, 1109, (astro-ph/0211536)
 Burwitz, V., Zavlin, V.E., Neuhaeuser, R., Predehl, P., Truemper, J., & Brinkman, A.C. 2001, A&A, 379, L35, (astro-ph/0109374)
 Caraveo, P.A., De Luca, A., Mereghetti, S., Pellizzoni, A., & Bignami, G.F. 2004, Science, 305, 376 (astro-ph/0407402)
 Chang, P., Morsink, M., Bildsten, L., & Wasserman, I. 2005, ApJ, 636, L117 (astro-ph/0511246)
 Cheng, K.S., Dai, Z.G., Wei, D.M., & Lu, T. 1998, Science, 280, 407
 Cook, G. B., Shapiro, S. L., Teukolsky, S. A. 1994, ApJ, 424, 823
 Cottam, J., Paerels, F., & Mendez, M. 2002, Nature, 420, 51
 Drake, J.J. et al. 2002, ApJ, 572, 996
 De Luca, A., Mereghetti, S., Caraveo, P.A., Moroni, M., Mignani R.P., Bignami, G.F. 2004, A&A, 418, 625
 Haensel, P. et al. 1986, A&A, 160, 121
 Haensel, P. 2001, A&A, 380, 186
 Homan, J., & van der Klis, M. 2000, ApJ, 539, 847 (astro-ph/0003241)
 Jansen, F. et al. 2001, A&A, 365, L1
 Jonker, P. G., van der Klis, M., & Groot, P. J. 2003, MNRAS, 339, 663
 Kaspi, V., Taylor, J., & Ryba, M. 1994, ApJ, 428, 713
 Lattimer, J. M., & Prakash, M. 2004, Science, 304, 536
 Lattimer, J. M., & Prakash, M. 2001, ApJ, 550, L426
 Li, X.D., Bombaci, I., Dey, M., Dey, J., & van den Heuvel, E.P.J. 1999, Phys. Rev. Lett., 3776, 83
 Lyne, A.G., Burgay, M., Kramer, M., Possenti, A., Manchester, R.N., Camilo, F., McLaughlin, M.A., Lorimer, D., D'Amico, N., Joshi, B.C., Reynolds, J., & Freire, P.C.C. 2004, Science, 303, 1089 (astro-ph/0401086)
 Makishima, K., Mihara, T., Nagase, F., & Tanaka, Y. 1999, ApJ, 525, 978
 Manchester, R. N., & Taylor, J. H. 1977, Pulsars (Freeman: San Francisco)
 Miller, M. C., Lamb, F. K., & Psaltis, D. 1998, ApJ, 508, 791
 Miller, M. C. 2002, Nature, 420, 31
 Morsink, S. 2000, Science, 290, 945
 Orosz, J.A., & Kuulkers, E. 1999, MNRAS, 305, 1320
 Ozel, F. 2006, Nature, 441, 1115 (astro-ph/0605106)
 Pavlov G. G., Sanwal D., Teter M. A. 2004, in Proceedings of Young Neutron Stars and Their Environments, IAU Symposium 218, Camilo F. and Gaensler B. M., eds., p.239 (astro-ph/0311526)
 Podsiadlowski, P., Dewi, J.D., Lesaffre, P., Miller, J.C., Newton, W.G., & Stone, J.R. 2005, "The Double Pulsar J0737-3039: Testing the Neutron Star Equation of State", (astro-ph/0506566)
 Rutledge, R. E., Bildsten, L., Brown, E. F., Pavlov, G. G., & Zavlin, V. E. 2001, ApJ, 559, 1054
 Sanwal, D., Pavlov, G. G., Zavlin, V. E., & Teter, M. A. 2002, ApJ, 574, L61
 Stella, L., & Vietri, M. 1999, Phys. Rev. Lett., 82, 17
 Thorne, K. S. 1977, ApJ, 212, 825
 Thorsett, S. E., & Chakrabarty, D. 1999, ApJ, 512, 288
 Tiengo, A., Galloway, D.K., di Salvo, T., Mendez, M., Miller, J.M., Sokoloski, J.L., & van der Klis, M. 2005, A&A accepted, (astro-ph/0506590)
 Truemper, J.E., Pietsch, W., Reppin, C. et al. 1978, ApJ, 219, L105
 Truemper, J.E., Burwitz, V., Haberl, F., & Zavlin, V.E. 2004, Nucl. Phys., 132, 156
 Truemper, J.E. 2005, private communication

- van der Klis, M. 2000, ARA&A, 38, 717 (astro-ph/0001167)
- van der Klis, M. 2006, "A review of rapid X-ray variability in X-ray binaries", to appear in Compact stellar X-ray sources, eds. W.H.G. Lewin & M. van der Klis, Cambridge University Press, p.39 (astro-ph/0410551)
- Villarreal, A.R., & Strohmayer, T.E. 2004, ApJ, 614, L121
- Walter, F.M., Lattimer, J. 2002, ApJ, 576, L145
- Xu, R. X. 2005, MNRAS, 356, 359
- Zhang, C.M. 2004, A&A, 423, 401 (astro-ph/0402028)
- Zhang, W., Smale, A.P., Strohmayer, T.E., & Swank, J.H. 1998, ApJ, 500, L171